

Fundamental Physical Constants — Physico-chemical constants

Quantity	Symbol	Value	Unit	Relative std. uncert. u_r
Avogadro constant	N_A, L	$6.022\,1415(10) \times 10^{23}$	mol^{-1}	1.7×10^{-7}
atomic mass constant $m_u = \frac{1}{12}m(^{12}\text{C}) = 1 \text{ u}$ $= 10^{-3} \text{ kg mol}^{-1}/N_A$	m_u	$1.660\,538\,86(28) \times 10^{-27}$	kg	1.7×10^{-7}
energy equivalent in MeV	$m_u c^2$	$1.492\,417\,90(26) \times 10^{-10}$ 931.494 043(80)	J MeV	1.7×10^{-7} 8.6×10^{-8}
Faraday constant ¹ $N_A e$	F	96 485.3383(83)	C mol^{-1}	8.6×10^{-8}
molar Planck constant	$N_A h$	$3.990\,312\,716(27) \times 10^{-10}$	J s mol^{-1}	6.7×10^{-9}
	$N_A hc$	0.119 626 565 72(80)	J m mol^{-1}	6.7×10^{-9}
molar gas constant	R	8.314 472(15)	$\text{J mol}^{-1} \text{ K}^{-1}$	1.7×10^{-6}
Boltzmann constant R/N_A in eV K ⁻¹	k	$1.380\,6505(24) \times 10^{-23}$ 8.617 343(15) $\times 10^{-5}$	J K^{-1} eV K ⁻¹	1.8×10^{-6} 1.8×10^{-6}
	k/h	2.083 6644(36) $\times 10^{10}$	Hz K ⁻¹	1.7×10^{-6}
	k/hc	69.503 56(12)	$\text{m}^{-1} \text{ K}^{-1}$	1.7×10^{-6}
molar volume of ideal gas RT/p $T = 273.15 \text{ K}, p = 101.325 \text{ kPa}$	V_m	$22.413\,996(39) \times 10^{-3}$	$\text{m}^3 \text{ mol}^{-1}$	1.7×10^{-6}
Loschmidt constant N_A/V_m	n_0	2.686 7773(47) $\times 10^{25}$	m^{-3}	1.8×10^{-6}
$T = 273.15 \text{ K}, p = 100 \text{ kPa}$	V_m	22.710 981(40) $\times 10^{-3}$	$\text{m}^3 \text{ mol}^{-1}$	1.7×10^{-6}
Sackur-Tetrode constant (absolute entropy constant) ² $\frac{5}{2} + \ln[(2\pi m_u k T_1/h^2)^{3/2} k T_1/p_0]$ $T_1 = 1 \text{ K}, p_0 = 100 \text{ kPa}$ $T_1 = 1 \text{ K}, p_0 = 101.325 \text{ kPa}$	S_0/R	-1.151 7047(44) -1.164 8677(44)		3.8×10^{-6} 3.8×10^{-6}
Stefan-Boltzmann constant $(\pi^2/60)k^4/\hbar^3c^2$	σ	$5.670\,400(40) \times 10^{-8}$	$\text{W m}^{-2} \text{ K}^{-4}$	7.0×10^{-6}
first radiation constant $2\pi hc^2$	c_1	$3.741\,771\,38(64) \times 10^{-16}$	W m^2	1.7×10^{-7}
first radiation constant for spectral radiance $2hc^2$	c_{1L}	$1.191\,042\,82(20) \times 10^{-16}$	$\text{W m}^2 \text{ sr}^{-1}$	1.7×10^{-7}
second radiation constant hc/k	c_2	$1.438\,7752(25) \times 10^{-2}$	m K	1.7×10^{-6}
Wien displacement law constant $b = \lambda_{\max}T = c_2/4.965\,114\,231\dots$	b	$2.897\,7685(51) \times 10^{-3}$	m K	1.7×10^{-6}

¹ The numerical value of F to be used in coulometric chemical measurements is 96 485.336(16) [1.7×10^{-7}] when the relevant current is measured in terms of representations of the volt and ohm based on the Josephson and quantum Hall effects and the internationally adopted conventional values of the Josephson and von Klitzing constants $K_{\text{J}-90}$ and $R_{\text{K}-90}$ given in the “Adopted values” table.

² The entropy of an ideal monoatomic gas of relative atomic mass A_r is given by $S = S_0 + \frac{3}{2}R \ln A_r - R \ln(p/p_0) + \frac{5}{2}R \ln(T/\text{K})$.